

## PHARMACY POLICY – 5.01.538


# ALK Tyrosine Kinase Inhibitors

Effective Date: July 1, 2024  
Last Revised: June 11, 2024  
Replaces: N/A

RELATED MEDICAL POLICIES:  
None

Select a hyperlink below to be directed to that section.

[POLICY CRITERIA](#) | [CODING](#) | [RELATED INFORMATION](#)  
[EVIDENCE REVIEW](#) | [REFERENCES](#) | [HISTORY](#)

 Clicking this icon returns you to the hyperlinks menu above.

---

## Introduction

The anaplastic lymphoma kinase (ALK) gene provides instructions for making a specific kind of protein called ALK receptor tyrosine kinase. This protein helps cells communicate. When this gene is damaged, cell growth can get stuck in the “on” position and cells grow uncontrollably. Changes to the ALK gene can lead to non-small-cell lung cancer. Tyrosine kinase inhibitors block specific enzymes, essentially working to turn the cell growth to the “off” position. ALK tyrosine kinase inhibitors specifically targets cancers caused by changes to the ALK gene. This policy describes when this specific form of chemotherapy may be considered medically necessary.

**Note:** The Introduction section is for your general knowledge and is not to be taken as policy coverage criteria. The rest of the policy uses specific words and concepts familiar to medical professionals. It is intended for providers. A provider can be a person, such as a doctor, nurse, psychologist, or dentist. A provider also can be a place where medical care is given, like a hospital, clinic, or lab. This policy informs them about when a service may be covered.

---

## Policy Coverage Criteria

Drug	Medical Necessity
<b>Alecensa (alectinib)</b>	<p><b>Alecensa (alectinib) may be considered medically necessary for the treatment of non-small cell lung cancer (NSCLC) when all the following are met:</b></p> <ul style="list-style-type: none"> <li>• The individual is 18 years of age or older</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual has been diagnosed with anaplastic lymphoma kinase (ALK)-positive NSCLC</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual meets one of the following: <ul style="list-style-type: none"> <li>○ Diagnosed with advanced or metastatic disease</li> <li>○ Requires adjuvant treatment following tumor resection with tumors <math>\geq</math> 4 cm or node positive</li> </ul> </li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The dose is limited to 1,200 mg daily</li> </ul>
<b>Alunbrig (brigatinib)</b>	<p><b>Alunbrig (brigatinib) may be considered medically necessary for the treatment of advanced or metastatic non-small cell lung cancer (NSCLC) when all the following are met:</b></p> <ul style="list-style-type: none"> <li>• The individual is 18 years of age or older</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual has been diagnosed with advanced or metastatic NSCLC that is anaplastic lymphoma kinase (ALK)-positive</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The dose is limited to <math>\leq</math> 180 mg daily</li> </ul> <p><b>Alunbrig (brigatinib) may be considered medically necessary for the treatment of inflammatory myofibroblastic tumor (iMT) with the ALK translocation.</b></p>
<b>Lorbrena (lorlatinib)</b>	<p><b>Lorbrena (lorlatinib) may be considered medically necessary for the treatment of advanced or metastatic non-small cell lung cancer (NSCLC) when all the following are met:</b></p> <ul style="list-style-type: none"> <li>• The individual is 18 years of age or older</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual has been diagnosed with advanced or metastatic NSCLC that is anaplastic lymphoma kinase (ALK)-positive</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The dose is limited to <math>\leq</math> 100 mg daily</li> </ul>



Drug	Medical Necessity
<b>Xalkori (crizotinib)</b>	<p><b>Xalkori (crizotinib) may be considered medically necessary for the treatment of advanced or metastatic non-small cell lung cancer (NSCLC) when all the following are met:</b></p> <ul style="list-style-type: none"> <li>• The individual is 18 years or older</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual has been diagnosed with advanced or metastatic NSCLC that is anaplastic lymphoma kinase (ALK)-positive or ROS proto-oncogene 1 receptor tyrosine kinase (ROS1)-positive</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The dose is limited to ≤ 500 mg daily</li> </ul> <p><b>Xalkori (crizotinib) may be considered medically necessary for the treatment of anaplastic large cell lymphoma (ALCL) when all the following are met:</b></p> <ul style="list-style-type: none"> <li>• The individual is between 1 year and 21 years of age</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual has been diagnosed with relapsed or refractory, systemic ALCL that is ALK-positive</li> </ul> <p><b>Xalkori (crizotinib) may be considered medically necessary for the treatment of inflammatory myofibroblastic tumor (iMT) when all the following are met:</b></p> <ul style="list-style-type: none"> <li>• The individual is 1 year of age or older</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual has been diagnosed with unresectable, recurrent, or refractory iMT that is ALK-positive</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The dose is limited to 500 mg daily</li> </ul>
<b>Zykadia (ceritinib)</b>	<p><b>Zykadia (ceritinib) may be considered medically necessary for the treatment of advanced or metastatic non-small cell lung cancer (NSCLC) when all the following are met:</b></p> <ul style="list-style-type: none"> <li>• The individual is 18 years of age or older</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>• The individual has been diagnosed with advanced or metastatic NSCLC that is anaplastic lymphoma kinase (ALK)-positive</li> </ul> <p><b>AND</b></p>



Drug	Medical Necessity
	<ul style="list-style-type: none"> <li>The dose is limited to <math>\leq 450</math> mg per daily</li> </ul> <p><b>Zykadia (ceritinib) may be considered medically necessary for the treatment of inflammatory myofibroblastic tumor (iMT) with the ALK translocation.</b></p>

Drug	Investigational
<b>As listed</b>	<b>All other uses of the medications listed in this policy are considered investigational.</b>

Length of Approval	
Approval	Criteria
<b>Initial authorization</b>	<b>All drugs listed in policy may be approved up to 3 months.</b>
<b>Re-authorization criteria</b>	<b>Future re-authorization of all drugs listed in policy may be approved up to 12 months as long as the drug-specific coverage criteria are met, and chart notes demonstrate that the individual continues to show a positive clinical response to therapy.</b>

Documentation Requirements
<p><b>The individual's medical records submitted for review for all conditions should document that medical necessity criteria are met. The record should include the following:</b></p> <ul style="list-style-type: none"> <li>Office visit notes that contain the diagnosis, relevant history, physical evaluation, and medication history</li> </ul>

## Coding

N/A



## Related Information

---

### Benefit Application

This policy is managed through the pharmacy benefit.

## Evidence Review

---

### Background

#### ALK+ Non-Small Cell Lung Cancer

Lung cancer consists of two major types: non-small cell lung cancer (NSCLC) and small-cell lung cancer (SCLC). Approximately 85% to 90% of lung cancers are NSCLC. NSCLC is further categorized into three major histological subtypes: adenocarcinoma, squamous cell (epidermoid) carcinoma, and large cell (undifferentiated) carcinoma. Adenocarcinoma is the most common subtype of NSCLC in the United States (about 40% of lung cancers). Several biomarkers have emerged as prognostic and predictive makers for NSCLC. These biomarkers include epidermal growth factor receptor (EGFR), the 5' endonuclease of the nucleotide excision repair complex (ERCC1), the k-ras oncogene, the regulatory subunit of ribonucleotide reductase (RRM1), and EML4-ALK fusion oncogene. Activation of ALK has been described as a primary oncogenic driver in about 2-7% of NSCLC individuals, or about 10,000 individuals in the United States. ALK-positive NSCLC is associated with distinct clinical features, including younger age of onset, absent or minimal smoking history, and adenocarcinoma histology.

In the United States, lung cancer is the second most common cancer and is the primary cause of cancer-related death in both men and women. It is estimated that 226,160 new cases of lung cancer (116,470 in men and 109,690 in women) are expected to be diagnosed in the United States in 2012. An estimated 160,340 deaths from lung cancer, accounting for about 28% of all cancer deaths, are expected to occur in the United States in 2012. More than 70% of individuals are diagnosed with stage III to IV disease. Survival outcomes in people with lung cancer vary depending on the stage of the cancer at diagnosis. Only about 15.6% of all lung cancer patients are alive 5 years or more after diagnosis. The financial burden of lung cancer treatment in the United States is estimated to be \$10.3 billion per year.



## Description

### The ALK Oncogene

Activation of ALK occurs through a chromosomal rearrangement that places one of several different 5' fusion partners and their associated promoter upstream of the 3' kinase domain of ALK. The most common 5' fusion partner in non-small cell lung cancer (NSCLC) is EML4, but other, rarer 5' fusion partners that cause oncogenic transformations have been described. The formation of ALK fusion proteins results in activation and dysregulation of gene expression and signaling which can contribute to increased cell proliferation and survival in tumors expressing these proteins. EML4-ALK translocations are usually mutually exclusive of epidermal growth factor receptor (EGFR) and KRAS mutations.

### Alecensa (alectinib)

Alecensa (alectinib) is a tyrosine kinase inhibitor indicated for the treatment of individuals with ALK-positive metastatic NSCLC. In nonclinical studies, alectinib inhibited ALK phosphorylation and ALK-mediated activation of the downstream signaling proteins STAT3 and AKT, and decreased tumor cell viability in multiple cell lines harboring ALK fusions, amplifications, or activating mutations. The major active metabolite of alectinib, M4, showed similar in vitro potency and activity. Alectinib and M4 demonstrated in vitro and in vivo activity against multiple mutant forms of the ALK enzyme, including some mutations identified in NSCLC tumors in individuals who have progressed on crizotinib.

### Alunbrig (brigatinib)

Alunbrig (brigatinib) has in vitro activity against multiple kinases including ALK, ROS1, IGF-1R, and FLT-3, as well as EGFR deletion and point mutations. Brigatinib inhibited autophosphorylation of ALK and ALK-mediated phosphorylation of the downstream signaling proteins STAT3, AKT, ERK1/2, and S6 in in vitro and in vivo assays. Brigatinib also inhibited the in vitro proliferation of cell lines expressing EML4-ALK and NPM-ALK fusion proteins. It inhibited the in vitro viability of cells expressing EML4-ALK and 17 mutant forms associated with resistance to ALK inhibitors EGFR-Del (E746-A750), ROS1-L2026M, FLT3-F691L, and FLT3-D835Y. Brigatinib exhibited in vivo anti-tumor activity against 4 mutant forms of EML4-ALK, including



G1202R and L1196M mutants identified in NSCLC tumors in individuals who have progressed on crizotinib.

## **Lorbrena (lorlatinib)**

Lorbrena (lorlatinib) received FDA approval in November 2018 for the treatment of ALK-positive metastatic NSCLC for individuals whose disease has progressed on other targeted therapies. In March 2021 Lorbrena was approved for first-line therapy based on results of the CROWN study. In this study patients were randomized 1:1 to receive lorlatinib 100 mg orally once daily or crizotinib 250 mg orally twice daily. Results from the study demonstrated a significant improvement in progression-free survival for the lorlatinib arm over the crizotinib arm. Lorlatinib has in vitro activity against ALK and ROS1 as well as TYK1, FER, FPS, TRKA, TRKB, TRKC, FAK, FAK2, and ACK. Lorlatinib demonstrated in vitro activity against multiple mutant forms of the ALK enzyme, including some mutations detected in tumors at the time of disease progression on crizotinib and other ALK inhibitors.

## **Xalkori (crizotinib)**

Xalkori (crizotinib) is a small-molecule tyrosine kinase receptor inhibitor, which inhibits ALK, Hepatocyte Growth Factor Receptor (HGFR, c-MET), and Recepteur d'Origine Nantais (RON). ALK gene abnormalities due to mutations or translocations may result in expression of oncogenic fusion proteins (i.e., EML4-ALK fusion protein) which alter signaling and expression and result in increased cellular proliferation and survival in tumors which express these fusion proteins. Crizotinib selectively inhibits ALK tyrosine kinase, which reduces proliferation of cells expressing the genetic alteration.

Xalkori (crizotinib) efficacy was demonstrated in two multicenter, single-arm studies that enrolled 255 patients with locally advanced or metastatic ALK-positive NSCLC: a phase II study (Study A [PROFILE 1005]) and a part two expansion cohort of a phase I dose-escalation study (Study B [Study 1001]). Patients enrolled into these studies had received prior systemic therapy, with the exception of 15 patients in Study B who had no prior systemic treatment for locally advanced or metastatic disease. The primary efficacy endpoint in both studies was investigator-determined Objective Response Rate (ORR) according to Response Evaluation Criteria in Solid Tumors (RECIST). Patients received 250 mg of crizotinib orally twice daily until disease progression or until intolerable side effects were reported. In Study A, there was 1 complete and 67 partial responses for an ORR of 50% (95% CI: 42%, 59%). In Study B, there were 2 complete



and 69 partial responses for an ORR of 61% (95% CI: 52%, 70%). The median response duration was 41.9 weeks and 48.1 weeks in Studies A and B, respectively.

Xalkori (crizotinib) was studied in individuals with ALK-positive inflammatory myofibroblastic tumors in two trials, ADVL0912 for pediatric patients and A8081013 for adults. The overall response rates were 86% (95% CI: 57%, 98%) for pediatric patients and 71% for adult patients.

Adverse effects were less than would be expected with conventional cytotoxic chemotherapy, but serious adverse events and treatment-related deaths related to crizotinib have been observed. Adverse events led to dosage reductions in 44% and 29% of patients in Studies A and B, respectively. Most adverse events were mild to moderate (Grade 1 or Grade 2). The most common ( $\geq 25\%$ ) adverse events were vision disorder, nausea, vomiting, diarrhea, edema, and constipation. Grade 3 or Grade 4 adverse events observed in  $\geq 2\%$  of patients included dyspnea, increased ALT levels, and neutropenia. Severe or fatal pneumonitis was reported in a small number of patients on crizotinib therapy.

## **Zykadia (ceritinib)**

Zykadia (ceritinib) is a small-molecule tyrosine kinase receptor inhibitor similar to crizotinib. Ceritinib was approved for treatment of ALK+ NSCLC on the basis of one single-arm, open-label, dose-finding Phase I study. This study started with 59 patients in the dose-escalation phase and recruited an additional 71 patients in the expansion phase, for a total of 130 patients. All patients had ALK-mutated tumors, with the majority (94%) with NSCLC; 68% of patients had previous progression of cancer while receiving crizotinib. The primary outcome was the finding the maximum tolerated dose and secondary outcome was tumor response as measured according to RECIST v1.0 by investigators and a blinded independent review committee (BIRC). The drop-out rate was not reported. There was a partial response in less than half of the patients (41.1% as evaluated by BIRC) and a complete response in only 2.5% of patients. The average progression free survival was 7 months. Previously approved second line agents (docetaxel, pemetrexed, erlotinib) did not have response rates greater than 10% and had progression free survival time of at most 12.3 weeks.

Common adverse events experienced with ceritinib include nausea, elevated liver function tests, diarrhea, and vomiting. Uncommon but serious events reported include interstitial lung disease, hyperglycemia, dyspnea, and prolonged QT interval. Fatal reactions occurred in 5% of patients in clinical trials due to pneumonia, respiratory failure, pneumonitis, pneumothorax, gastric hemorrhage, pulmonary tuberculosis, cardiac tamponade, and sepsis.





## **2013 Update**

Added treatment of inflammatory myofibroblastic tumor with ALK translocation (NCCN category 2A).

## **2014 Update**

Policy updated with new ALK tyrosine kinase inhibitor, ceritinib.

## **2015 Update**

Added treatment of ROS1-positive non-small-cell lung cancer with crizotinib (NCCN category 2A).

## **2016 Update**

Policy updated with new ALK tyrosine kinase inhibitor, alectinib.

## **2017 Update**

Policy updated with literature review for the previous year. A statement outlining the length of therapy for initial and subsequent approval has been added to the policy.

## **2018 Update**

Policy updated with no changes. Third-generation inhibitor loratinib is currently in phase III clinical trials, expecting FDA approval in August 2018. Added reauthorization criteria and documentation statement.



## 2019 Update

Reviewed prescribing information for all drugs. No new evidence was identified that would require changes to drugs listed in this policy.

## 2020 Update

Reviewed prescribing information for all drugs. No new evidence was identified that would require changes to drugs listed in this policy. Added a Documentation Requirements table to policy.

## 2021 Update

Reviewed prescribing information for all drugs. Based on results from the CROWN study in previously untreated ALK-positive NSCLC, Lorbrena (lorlatinib) coverage criteria were updated removing requirement for prior treatment with an alternative ALK kinase inhibitor. Coverage criteria for Alecensa (alectinib), Alunbrig (brigatinib), Xalkori (crizotinib), and Zykadia (ceritinib) for the treatment of NSCLC were updated limiting use to adult individuals. Quantity limits were also added to all drugs for the treatment of NSCLC following the prescribing information. Per the NCCN Guidelines Version 2.2021 for Soft Tissue Sarcoma, for the treatment of inflammatory myofibroblastic tumor (iMT) with the ALK translocation Alunbrig (brigatinib), Xalkori (crizotinib), and Zykadia (ceritinib) are all listed as preferred ALK inhibitors with a category 2A recommendation. Based on NCCN Guidelines, coverage criteria were expanded for Alunbrig (brigatinib) and Zykadia (ceritinib) to include treatment of iMT with the ALK translocation. References and background information were revised.

## 2022 Update

Reviewed prescribing information for all drugs. Reviewed new indication for Xalkori (crizotinib) and related pivotal trials. Added coverage for Xalkori (crizotinib) for ALK-positive inflammatory myofibroblastic tumors.



## 2023 Update

Reviewed prescribing information for all drugs in the policy. No new evidence was found that could change the policy statements. Changed the wording from "patient" to "individual" throughout the policy for standardization.

## 2024 Update

Reviewed prescribing information for all drugs in the policy. Updated Alecensa (alectinib) coverage criteria to include treatment of certain adults requiring adjuvant treatment for non-small cell lung cancer (NSCLC).

## References

1. Kwak EL, Bang, YJ, Camidge DR. Anaplastic lymphoma kinase inhibition in non-small cell lung cancer. *N Eng J Med*. 2010;363(18):1693-1703.
2. Shepherd FA, Dancey J, Ramlau R, et al. Prospective randomized trial of docetaxel versus best supportive care in patients with non-small-cell lung cancer previously treated with platinum-based chemotherapy. *J Clin Oncol*. 2000;18(10):2095–2103.
3. ClinicalTrials.gov. An Investigational Drug, PF-02341066, Is Being Studied In Patients With Advanced Non-Small Cell Lung Cancer With A Specific Gene Profile Involving The Anaplastic Lymphoma Kinase (ALK) Gene (NCT00932451). Available at: <http://clinicaltrials.gov/ct2/show/NCT00932451?term=NCT00932451&rank=1> Accessed June 4, 2024.
4. Center for Drug Evaluation and Research. Medical Review(s) application number: 20257Orig1s000. March 30, 2011. Available at: [http://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2011/202570Orig1s000TOC.cfm](http://www.accessdata.fda.gov/drugsatfda_docs/nda/2011/202570Orig1s000TOC.cfm) Accessed June 4, 2024.
5. Xalkori (crizotinib) product dossier. Pfizer, Inc; New York, NY. Revised September 2023.
6. Center for Drug Evaluation and Research. Statistical Review(s)-application number: 20257Orig1s000. March 30, 2011. Available at: [http://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2011/202570Orig1s000TOC.cfm](http://www.accessdata.fda.gov/drugsatfda_docs/nda/2011/202570Orig1s000TOC.cfm) Accessed June 4, 2024.
7. Center for Drug Evaluation and Research. Cross Discipline Team Leader Review-application number: 20257Orig1s000. July 12, 2010. Available at: [http://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2011/202570Orig1s000TOC.cfm](http://www.accessdata.fda.gov/drugsatfda_docs/nda/2011/202570Orig1s000TOC.cfm) Accessed June 4, 2024.
8. ClinicalTrials.gov. Phase 3, Randomized, Open-Label Study Of The Efficacy And Safety Of PF-02341066 Versus Standard Of Care Chemotherapy (Pemetrexed Or Docetaxel) In Patients With Non-Small Lung Cancer Harboring A Translocation Or Inversion Event Involving The Anaplastic Lymphoma Kinase (ALK) Gene Locus (NCT00932893). Available at: <http://clinicaltrials.gov/ct2/show/NCT00932893> Accessed June 4, 2024.
9. ClinicalTrials.gov. Phase 3, Randomized, Open-Label Study Of The Efficacy And Safety Of Crizotinib Versus Pemetrexed/Cisplatin Or Pemetrexed/Carboplatin In Previously Untreated Patients With Non-Squamous Carcinoma Of The Lung Harboring A Translocation Or Inversion Event Involving The Anaplastic Lymphoma Kinase (ALK) Gene Locus (NCT01154140). Available at: <https://clinicaltrials.gov/ct2/show/NCT01154140> Accessed June 4, 2024.



10. National Comprehensive Cancer Network (NCCN) Guidelines: Soft Tissue Sarcoma, Version 3.2021. Available at [https://www.nccn.org/professionals/physician\\_gls/pdf/sarcoma.pdf](https://www.nccn.org/professionals/physician_gls/pdf/sarcoma.pdf) Accessed June 4, 2024.
11. Azzoli CG, Temin S, Aliff T, et al. 2011 focused update of 2009 American Society of Clinical Oncology clinical practice guideline update on chemotherapy for stage IV non–small cell lung cancer. *J Clin Oncol*. 2011;29(28):3825-3831.
12. Zykadia (ceritinib) Prescribing Information. Novartis Pharmaceuticals, East Hanover, NJ. Revised October 2021.
13. Shaw AT, Kim DW, Mehra R, et al. Ceritinib in ALK-rearranged non-small-cell lung cancer. *N Engl J Med*. Mar 27 2014;370(13):1189-1197.
14. ClinicalTrials.gov. A Phase I, Multi-center, Open Label Dose Escalation Study of LDK378, Administered Orally in Adult Patients With Tumors Characterized by Genetic Abnormalities in Anaplastic Lymphoma Kinase (ALK). Available at: <http://clinicaltrials.gov/show/NCT01283516>. Accessed June 4, 2024.
15. Shaw AT, Ou SI, Bang YJ, et al. Crizotinib in ROS1-Rearranged Non-Small-Cell Lung Cancer. *N Engl J Med*. Nov 20 2014;371:1963-71.
16. Alecensa (alectinib) Prescribing Information. Genentech USA, Inc., South San Francisco, CA. Revised April 2024.
17. Alunbrig (brigatinib) Prescribing Information. Takeda Pharmaceutical Company Limited, Cambridge, MA. Revised February 2022.
18. Lorbreña (lorlatinib) Prescribing Information. Pfizer Inc., New York, New York. Updated April 2023.

## History

Date	Comments
06/12/12	New policy, add to Prescription Drug section.
07/08/13	Minor Update – Clarification was added to the policy that it is managed through the member’s pharmacy benefit; this is now listed in the header and within the coding section.
10/14/13	Replace policy. Medically necessary indications for crizotinib expanded to include treatment of inflammatory myofibroblastic tumor (iMT) with the ALK translocation.
11/10/14	Annual Review. Policy updated with literature review; medically necessary policy statement added for the new ALK tyrosine kinase inhibitor, ceritinib. References 12-15 added.
09/08/15	Annual Review. Policy updated with literature review; reference 16 added. Medically necessary policy statement for Xalkori (crizotinib) updated to include the indication of ROS proto-oncogene 1 receptor tyrosine kinase (ROS1)-positive.
04/01/16	Interim update, changes approved March 8, 2016. Medically necessary policy statement added for the new ALK tyrosine kinase inhibitor, alectinib.
05/01/17	Annual Review, changes approved April 11, 2017. A statement outlining the length of therapy for initial and subsequent approval has been added to the policy.
07/01/17	Interim update. approved June 13, 2017. Policy moved into the new format. Added coverage criteria for Alunbrig (brigatinib).



Date	Comments
10/01/17	Interim review approved September 21, 2017. Alecensa, Zykadia, Alunbrig changed to first-line.
07/01/18	Annual Review, approved June 22, 2018. Added reauthorization criteria and documentation statement.
03/01/19	Interim Review, approved February 12, 2019. Added coverage criteria for Lorbrena (lorlatinib).
12/01/19	Annual Review, approved November 21, 2019. Literature review through October 2019; no change to policy statements.
01/01/21	Annual Review, approved December 1, 2020. No changes to policy statements.
03/01/21	Interim Review, approved February 18, 2021. Added a new indication to Xalkori (crizotinib) for the treatment of anaplastic large cell lymphoma that is ALK-positive.
06/01/21	Annual Review, approved May 20, 2021. Updated coverage criteria for Lorbrena (lorlatinib) for the treatment of NSCLC removing requirement for prior treatment with an ALK kinase inhibitor and adding a daily dose limit. Updated coverage criteria for Alecensa (alectinib), Alunbrig (brigatinib), Xalkori (crizotinib), and Zykadia (ceritinib) for the treatment of NSCLC limiting use to adults and adding a daily dose limit. Expanded coverage for Alunbrig (brigatinib) and Zykadia (ceritinib) to include treatment of inflammatory myofibroblastic tumor (iMT) with the ALK translocation.
04/01/22	Annual Review, approved March 7, 2022. No changes to policy statements.
10/01/22	Interim Review, approved September 13, 2022. Updated coverage for Xalkori (crizotinib) for ALK-positive inflammatory myofibroblastic tumors (IMT) to limit use to patients 1 year of age and older with unresectable, recurrent, or refractory IMT.
09/01/23	Annual Review, approved August 7, 2023. Reviewed prescribing information for all drugs in the policy. No new evidence was found that could change the policy statements. Changed the wording from "patient" to "individual" throughout the policy for standardization.
07/01/24	Annual Review, approved June 11, 2024. Updated Alecensa (alectinib) coverage criteria to include treatment of certain adults requiring adjuvant treatment for non-small cell lung cancer (NSCLC).

**Disclaimer:** This medical policy is a guide in evaluating the medical necessity of a particular service or treatment. The Company adopts policies after careful review of published peer-reviewed scientific literature, national guidelines and local standards of practice. Since medical technology is constantly changing, the Company reserves the right to review and update policies as appropriate. Member contracts differ in their benefits. Always consult the member benefit booklet or contact a member service representative to determine coverage for a specific medical service or supply. CPT codes, descriptions and materials are copyrighted by the American Medical Association (AMA). ©2024 Premera All Rights Reserved.



**Scope:** Medical policies are systematically developed guidelines that serve as a resource for Company staff when determining coverage for specific medical procedures, drugs or devices. Coverage for medical services is subject to the limits and conditions of the member benefit plan. Members and their providers should consult the member benefit booklet or contact a customer service representative to determine whether there are any benefit limitations applicable to this service or supply. This medical policy does not apply to Medicare Advantage.



**Discrimination is Against the Law**

Premera Blue Cross HMO (Premera HMO) complies with applicable Federal and Washington state civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, sex, gender identity, or sexual orientation. Premera HMO does not exclude people or treat them differently because of race, color, national origin, age, disability, sex, gender identity, or sexual orientation. Premera HMO provides free aids and services to people with disabilities to communicate effectively with us, such as qualified sign language interpreters and written information in other formats (large print, audio, accessible electronic formats, other formats). Premera HMO provides free language services to people whose primary language is not English, such as qualified interpreters and information written in other languages. If you need these services, contact the Civil Rights Coordinator. If you believe that Premera HMO has failed to provide these services or discriminated in another way on the basis of race, color, national origin, age, disability, sex, gender identity, or sexual orientation, you can file a grievance with: Civil Rights Coordinator — Complaints and Appeals, PO Box 91102, Seattle, WA 98111, Toll free: 855-332-4535, Fax: 425-918-5592, TTY: 711, Email [AppealsDepartmentInquiries@Premera.com](mailto:AppealsDepartmentInquiries@Premera.com). You can file a grievance in person or by mail, fax, or email. If you need help filing a grievance, the Civil Rights Coordinator is available to help you. You can also file a civil rights complaint with the U.S. Department of Health and Human Services, Office for Civil Rights, electronically through the Office for Civil Rights Complaint Portal, available at <https://ocrportal.hhs.gov/ocr/portal/lobby.jsf>, or by mail or phone at: U.S. Department of Health and Human Services, 200 Independence Ave SW, Room 509F, HHH Building, Washington, D.C. 20201, 1-800-368-1019, 800-537-7697 (TDD). Complaint forms are available at <http://www.hhs.gov/ocr/office/file/index.html>. You can also file a civil rights complaint with the Washington State Office of the Insurance Commissioner, electronically through the Office of the Insurance Commissioner Complaint Portal available at <https://www.insurance.wa.gov/file-complaint-or-check-your-complaint-status>, or by phone at 800-562-6900, 360-586-0241 (TDD). Complaint forms are available at <https://fortress.wa.gov/oic/onlineservices/cc/pub/complaintinformation.aspx>.

**Language Assistance**

- ATENCIÓN:** si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 844-722-4661 (TTY: 711).
- 注意:** 如果您使用繁體中文，您可以免費獲得語言援助服務。請致電 844-722-4661 (TTY: 711)。
- CHÚ Ý:** Nếu bạn nói Tiếng Việt, có các dịch vụ hỗ trợ ngôn ngữ miễn phí dành cho bạn. Gọi số 844-722-4661 (TTY: 711).
- 주의:** 한국어를 사용하시는 경우, 언어 지원 서비스를 무료로 이용하실 수 있습니다. 844-722-4661 (TTY: 711) 번으로 전화해 주십시오.
- ВНИМАНИЕ:** Если вы говорите на русском языке, то вам доступны бесплатные услуги перевода. Звоните 844-722-4661 (телетайп: 711).
- PAUNAWA:** Kung nagsasalita ka ng Tagalog, maaari kang gumamit ng mga serbisyo ng tulong sa wika nang walang bayad. Tumawag sa 844-722-4661 (TTY: 711).
- УВАГА!** Якщо ви розмовляєте українською мовою, ви можете звернутися до безкоштовної служби мовної підтримки. Телефонуйте за номером 844-722-4661 (телетайп: 711).
- ប្រយ័ត្ន:** បើសិនជាអ្នកនិយាយ ភាសាខ្មែរ, សេវាជំនួយផ្នែកភាសា ដោយមិនគិតលុយ គឺអាចមានសំរាប់អ្នក។ ចូរ ទូរស័ព្ទ 844-722-4661 (TTY: 711)។
- 注意事項:** 日本語を話される場合、無料の言語支援をご利用いただけます。844-722-4661 (TTY:711) まで、お電話にてご連絡ください。
- ማስታወሻ:** የሚናገሩት ቋንቋ አማርኛ ከሆነ የትርጉም እርዳታ ድርጅቶች፣ በገጻ ሊያግዝዎት ተዘጋጅተዋል። ወደ ሚከተለው ቁጥር ይደውሉ 844-722-4661 (መስማት ለተሳናቸው: 711)።
- XIYYEEFFANNA:** Afaan dubbattu Oroomiffa, tajaajjila gargaarsa afaanii, kanfaltiidhaan ala, ni argama. Bilbilaa 844-722-4661 (TTY: 711).
- ملحوظة:** إذا كنت تتحدث اذكر اللغة، فإن خدمات المساعدة اللغوية تتوافر لك بالمجان. اتصل برقم 844-722-4661 (رقم هاتف الصم والبكم: 711).
- ਧਿਆਨ ਦਿਓ:** ਜੇ ਤੁਸੀਂ ਪੰਜਾਬੀ ਬੋਲਦੇ ਹੋ, ਤਾਂ ਭਾਸ਼ਾ ਵਿੱਚ ਸਹਾਇਤਾ ਸੇਵਾ ਤੁਹਾਡੇ ਲਈ ਮੁਫਤ ਉਪਲਬਧ ਹੈ। 844-722-4661 (TTY: 711) 'ਤੇ ਕਾਲ ਕਰੋ।
- ACHTUNG:** Wenn Sie Deutsch sprechen, stehen Ihnen kostenlos sprachliche Hilfsdienstleistungen zur Verfügung. Rufnummer: 844-722-4661 (TTY: 711).
- ໂປດອຸບ:** ຖ້າວ່າ ທ່ານວົ້າພາສາ ລາວ, ການບໍລິການຊ່ວຍເຫຼືອດ້ານພາສາ, ໂດຍບໍ່ຄ່າສ່ຽງຄ່າ, ຄວນມີພ້ອມໃຫ້ທ່ານ. ໂທ 844-722-4661 (TTY: 711).
- ATANSYON:** Si w pale Kreyòl Ayisyen, gen sèvis èd pou lang ki disponib gratis pou ou. Rele 844-722-4661 (TTY: 711).
- ATTENTION:** Si vous parlez français, des services d'aide linguistique vous sont proposés gratuitement. Appelez le 844-722-4661 (ATS : 711).
- UWAGA:** Jeżeli mówisz po polsku, możesz skorzystać z bezpłatnej pomocy językowej. Zadzwoń pod numer 844-722-4661 (TTY: 711).
- ATENÇÃO:** Se fala português, encontram-se disponíveis serviços linguísticos, grátis. Ligue para 844-722-4661 (TTY: 711).
- ATTENZIONE:** In caso la lingua parlata sia l'italiano, sono disponibili servizi di assistenza linguistica gratuiti. Chiamare il numero 844-722-4661 (TTY: 711).
- توجه:** اگر بہ زبان فارسی گفتگو می کنید، تسهیلات زبانی بصورت رایگان برای شما فراهم می باشد. با 844-722-4661 (TTY: 711) تماس بگیرید.